

Citation for published version:

Turner, JE, Wadley, A, Aldred, S, Fisher, J, Bosch, JA & Campbell, JP 2016, 'Intensive exercise does not preferentially mobilise skin-homing T cells and NK cells', *Medicine & Science in Sports & Exercise*, vol. 48, no. 7, pp. 1285-1293. <https://doi.org/10.1249/MSS.0000000000000914>

DOI:

[10.1249/MSS.0000000000000914](https://doi.org/10.1249/MSS.0000000000000914)

Publication date:

2016

Document Version

Peer reviewed version

[Link to publication](#)

This is the author accepted manuscript of an article published in final form as TURNER, JAMES E.; WADLEY, ALEX J.; ALDRED, SARAH; FISHER, JAMES P.; BOSCH, JOS A.; CAMPBELL, JOHN P. Intensive Exercise Does Not Preferentially Mobilize Skin-Homing T Cells and NK Cells, *Medicine & Science in Sports & Exercise*: July 2016 - Volume 48 - Issue 7 - p 1285-1293 and available online via: <https://doi.org/10.1249/MSS.0000000000000914>

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Intensive exercise does not preferentially mobilise skin-homing T cells and NK cells.**

2 **Running head:** CLA+ lymphocytes and exercise

3

4 ¹James E. Turner, ^{2,3}Alex J. Wadley, ³Sarah Aldred, ³James P. Fisher, ^{4,5}Jos A. Bosch, ^{*6}John P. Campbell.

5

6 1. Department for Health, University of Bath, Bath, UK.

7 2. Institute of Science and the Environment, University of Worcester, Worcester, UK.

8 3. School of Sport, Exercise and Rehabilitation Sciences, College of Life and Environmental Sciences,
9 University of Birmingham, Birmingham, UK.

10 4. Department of Psychology, University of Amsterdam, Amsterdam, The Netherlands.

11 5. Mannheim Institute of Public Health, Social and Preventive Medicine (MIPH), University of Heidelberg,
12 Heidelberg, Germany

13 6. Institute of Immunology and Immunotherapy, University of Birmingham, Birmingham, UK.

14

15 * Corresponding author details:

16 John P. Campbell

17 Institute of Immunology and Immunotherapy, University of Birmingham, Birmingham, UK. B15 2TT.

18 **Telephone:** +44 7530595912

19 **Email:** j.p.campbell@bham.ac.uk

20

21

22 **Abstract**

23 **Purpose:** This study investigated whether natural killer (NK) cells and CD8⁺ T cells expressing cutaneous
24 lymphocyte antigen (CLA) – a homing molecule for endothelial cell leukocyte adhesion molecule 1 (ELAM-
25 1), which enables transmigration to the skin – are selectively mobilised in response to acute exercise.
26 **Methods:** Nine healthy males (mean \pm SD age: 22.1 \pm 3.4 years) completed two exercise sessions: high-
27 intensity continuous cycling ('continuous exercise' at 80% $\dot{V}O_{2\text{MAX}}$ for 20 min) and low-volume high-
28 intensity interval exercise ('HIIE' at 90% $\dot{V}O_{2\text{MAX}}$ 10 \times 1 min repetitions with 1 min recovery intervals).
29 Blood was collected before, immediately- and 30 min post-exercise for cryo-preservation of peripheral blood
30 mononuclear cells. CLA⁺ and CLA[−] cells were quantified within NK subpopulations (CD56^{bright} 'regulatory'
31 and CD56^{dim} 'cytotoxic' cells) as well as the following CD8⁺ T cell subpopulations: naive ('NA';
32 CD45RA+CCR7+), central memory ('CM'; CD45RA−CCR7+), effector-memory ('EM'; CD45RA−CCR7−)
33 and CD45RA-expressing effector-memory cells ('EMRA'; CD45RA+CCR7−). **Results:** CLA⁺ NK cells and
34 CD8⁺ memory T cells increased in response to both exercise bouts, but, overall, their numerical contribution
35 to the exercise lymphocytosis was inferior to CLA[−] cells, which increased to a much greater extent during
36 exercise. Tellingly, the most exercise-responsive cells – effector memory CD8⁺ cells and CD56^{dim} cells –
37 were CLA[−]. **Conclusions:** A small subset of CLA⁺ lymphocytes are mobilised into blood during acute
38 intensive exercise, but CLA⁺ cells are not major contributors to exercise lymphocytosis, thus providing
39 preliminary evidence that the skin is not a major origin, or homing-destination, of exercise-sensitive
40 lymphocytes.

41

42 **Key words:** T cells, inflammation, skin, homing

43

44 INTRODUCTION

45 Memory CD8⁺ T cells and natural killer (NK) cells are the largest contributors to the acute and
46 transient lymphocytosis that occurs in peripheral blood during intensive exercise (6, 20, 36). Largely driven
47 by their relative high density cell surface expression of β 2-adrenergic receptors that induce detachment from
48 the vascular endothelium upon stimulation by catecholamines, the mobilisation of NK cell and CD8⁺ T cell
49 subsets is considered to be an evolved mechanism that facilitates effector cell recruitment to sites of potential
50 or ongoing injury (10). Built on findings from animal models and using an analogy of immune cells as
51 ‘soldiers’, it is proposed that naïve and central memory CD8⁺ T cells – which are slightly increased during
52 exercise – traffic from the boulevards (i.e., ‘bloodstream’) back to the ‘barracks’ (i.e., the spleen, lymph
53 nodes). On the other hand, tissue-specific memory CD8⁺ T cells – which mobilise to a larger extent during
54 exercise – home to specific peripheral tissues (e.g., lungs, gut) to conduct immune-surveillance against
55 potential pathogenic challenge. Finally, it is hypothesised that effector CD8⁺ T cells – which are mobilised to
56 the greatest extent during exercise – are redeployed to ‘battlefield’ sites of wound healing (e.g. in the skin)
57 following acute exercise (9, 10). However, whether these highly cytotoxic effector NK and CD8⁺ T cells,
58 which are selectively increased in an acute and transient manner by acute exercise, have the propensity to
59 migrate to peripheral cutaneous sites remains unknown.

60 In support of this model, it has been demonstrated in murine models that exercise redeploys large
61 numbers of T cells to the Peyer’s patches, lungs and bone marrow, reinforcing the idea that T cells are
62 mobilised to sites of potential antigen encounter (i.e., lungs, gut), as well as to serve other important functions
63 such as the provision of additional stimuli for haematopoiesis in the bone marrow (20). However, the
64 aforementioned study was unable to measure homing to cutaneous sites or areas of ongoing inflammation. It
65 has been shown that T cells and NK cells do show increased infiltration into sites of experimental
66 inflammation (i.e., subcutaneous implantation of a surgical sponge treated with the lymphocyte-specific
67 chemokine lymphotactin) in mice 24 to 48 hours after acute psychological stress (40). However, it remains
68 unclear whether this increased cell infiltration is driven by stress-induced effector cell redistribution, or other
69 mechanisms occurring in the days after the acute stressor.

70 An approach that is commonly used to investigate lymphocyte homing propensity in humans is the
71 assessment of cell surface adhesion molecule expression on cells; this approach can be used to reveal the
72 probable trafficking patterns of cells mobilised into the bloodstream during exercise. For example, studies
73 have shown that there is a selective influx of CD8⁺ memory cells into the bloodstream that exhibit lower
74 levels of lymphoid homing markers such as CD62L and CCR7 (6), thus providing evidence that exercise
75 mobilises CD8⁺ T cells with a homing capacity for peripheral tissues. Further research has shown that these
76 cells mobilised by exercise express adhesion molecules such as CD11a (15, 22), CD11b (17, 18), VLA-4
77 (very late antigen-4) and LPAM-1 (lymphocyte Peyer's patch adhesion molecule-1) (15), which enable
78 migration to peripheral sites including the bone marrow (23), Peyer's patches (43) and lungs (38). However,
79 the aforementioned adhesion molecules cannot be used exclusively for identifying skin-homing potential and
80 it remains uncertain whether exercise-responsive CD8⁺ T cells and NK cells have a skin-homing phenotype.

81 Cutaneous lymphocyte antigen (CLA) expression on lymphocytes can be used to determine whether
82 lymphocytes preferentially mobilised during intensive exercise exhibit a homing phenotype for cutaneous
83 sites. CLA is a specialised form of P-selectin glycoprotein ligand-1 (PSGL-1; CD162), a surface glycoprotein
84 expressed constitutively on all human peripheral-blood T cells. After post-translational modification, PSGL-1
85 bears a Sialyl-Lewis^X (sLe^X) moiety, termed CLA, which avidly binds CD62E (E-selectin), an adhesion
86 molecule which initiates the transmigration cascade to the skin, and which is also upregulated during
87 cutaneous inflammation (4, 27, 28, 32). Thus, most T cells in both normal and diseased cutaneous sites are
88 CLA⁺ (7, 29, 30). In support, leukocyte infiltration to inflammatory sites can be largely inhibited by a CLA
89 modifier (11). Taken together, analyses of CLA⁺ cell mobilisation can be used as proxy marker to reveal
90 whether cells mobilised by exercise have the phenotypic capacity to migrate to 'battlefield' sites in the skin
91 (10).

92 The primary objective of this study was to investigate whether CD8⁺ T cells preferentially mobilised
93 during intensive exercise stress exhibit a homing phenotype for cutaneous sites. To fulfil this objective, we
94 investigated the number of CLA-positive cells among CD8⁺ T cell subsets, to establish the contribution of
95 CLA-positive cells to the stepwise CD8⁺ lymphocytosis pattern previously observed in response to exercise
96 (6). We also extended the analyses of CLA⁺ and CLA⁻ cells to NK cell populations – the largest responders

97 to exercise stress (6). The second objective of this study was to compare the magnitude of CD8+ T cell and
98 NK cell mobilisation in response to continuous high-intensity exercise (continuous exercise) and high-
99 intensity intermittent exercise (HIIE). HIIE is often referred to as High Intensity Interval Training (HIIT)
100 when repeated frequently over several weeks or months. This form of exercise has received considerable
101 attention as a short-duration and low-volume means of achieving similar health benefits to continuous
102 exercise. Given the increasing health benefits that have been observed with HIIE, and although we have
103 established in a prior report that CD8+ T cell and NK cell subset mobilisation is intensity dependent (6), the
104 effects of HIIE on changes to these cell subsets remains unknown.

METHODS

Participants

Ten healthy males were recruited to take part in this study as previously described (41, 42). Peripheral blood mononuclear cells (PBMCs) were available from nine of the ten participants, and were isolated from blood samples collected before, immediately after, and 30 min after two different forms of exercise, described below. All nine participants (age: 22.1 ± 3.4 years; height: 180.5 ± 6.1 cm; weight: 78.1 ± 11.0 kg; body mass index: 24.0 ± 3.1 kg.m²; $\dot{V}O_{2\text{MAX}}$: 43.8 ± 4.1 ml.kg.min⁻¹) included in this study were non-smokers, and refrained from taking vitamin supplements and anti-inflammatory medication for fourteen days, and did not exercise and consume alcohol or caffeine for two days prior to experimental trials. All participants provided written informed consent and the study was approved by the Science, Technology, Engineering and Mathematics Ethical Review Committee at University of Birmingham (reference: ERN_12-0830).

Preliminary measurements

Height and weight were assessed using standard methods and cardiorespiratory fitness ($\dot{V}O_{2\text{MAX}}$) was measured during an incremental exercise test on an electromagnetically braked cycle ergometer (*Lode Excalibur Sport*, Groningen, Netherlands). Following a three-minute warm up at 30 watts, workload was increased by 30 watts every minute, until volitional exhaustion. A facemask was fitted throughout preliminary exercise tests in each main trial, and expired air measured breath-by-breath averaged every 20 seconds for oxygen uptake and carbon dioxide production (*Oxycon Pro*, Jaeger, Wuerzburg, Germany). Heart rate (HR) was monitored every minute using a Polar Vantage heart rate monitor (Polar Vantage, Kempele, Finland); HR_{peak} represents the maximum heart rate achieved during each trial. The following criteria were used to indicate that $\dot{V}O_{2\text{MAX}}$ had been reached: a fall in cadence below 60 rpm, a respiratory exchange ratio ($\dot{V}CO_2 / \dot{V}O_2$) >1.10-1.15, a plateau in oxygen consumption and a heart rate >220 beats min⁻¹ minus age.

Experimental trials

Experimental trials were undertaken at least seven days after preliminary measurements, in the morning, and following an overnight fast. Each trial was separated by at least three days in a randomised

design. Prior to each trial, but after the baseline blood sample, participants undertook a warm up (5 minutes) at a workload eliciting 40% $\dot{V}O_{2\text{MAX}}$. Exercise trials were either vigorous steady state cycling at 80% $\dot{V}O_{2\text{MAX}}$ for 20 minutes ('continuous exercise') or high intensity interval exercise ('HIIE') trial. HIIE consisted of ten 1 minute cycling phases at a workload to elicit 90% $\dot{V}O_{2\text{MAX}}$, with 1 minute of low intensity cycling at 40% $\dot{V}O_{2\text{MAX}}$ between each phase. Workload was expressed as watts, and relative to body mass (i.e., watts/kg). Values presented are the average workload over the entire exercise protocol (i.e., in the vigorous trial: the duration of cycling at 80% $\dot{V}O_{2\text{MAX}}$; and in the HIIE protocol: each 1-minute sprint phase of cycling at 90% $\dot{V}O_{2\text{MAX}}$). Values were obtained directly from the electromagnetically braked cycle ergometer (Lode Excalibur Sport, Groningen, Netherlands). Energy expenditure (Kcal) was estimated by indirect calorimetry from calculations of fat and carbohydrate oxidation (g/min) and data were averaged over the 1-minute phases of the protocols and summed to provide total energy expenditure from the entire duration of each trial, expressed relative to body mass (Kcal/kg). Ratings of perceived exertion (RPE) were recorded using the Borg Scale every 1 minute during continuous exercise, and after each 90% $\dot{V}O_{2\text{MAX}}$ interval during HIIE; the highest RPE score was selected as the final RPE result for each exercise trial (RPE_{peak}).

Blood sampling

An intravenous cannula (*Becton & Dickson, Oxford, UK*) was inserted into an antecubital vein and blood samples were drawn into potassium ethylene diaminetetraacetic acid (EDTA) vacutainer tubes (*Becton & Dickson, Oxford, UK*). The cannula was kept patent with saline (0.9% NaCl). The leukocyte differential was assessed using an automated haematology analyser (*Coulter Analyser, Beckman-Coulter, High Wycombe, UK*).

Blood Cell Isolation

Approximately 15 ml of blood from each time point (pre-exercise, post-exercise and 30 min post-exercise) was diluted 1:1 with Roswell Park Memorial Institute Media (RPMI), and then layered on top of Ficoll paque PLUS (GE Healthcare) (2 blood : 1 Ficoll), before centrifuging at $500 \times g$ for 30 minutes at 21°C. PBMCs were aspirated and washed three times in RPMI by centrifuging at $400 \times g$ for 5 minutes. The

cell pellet was re-suspended in 1-ml of freezing mixture (70% RPMI, 20% fetal calf serum (FCS) and 10% dimethyl sulfoxide (DMSO)) and frozen at -1°C / min using a freezing container (Nalgene “Mr Frosty” Thermoscientific). Cells were stored at -80°C and analysed within six months.

162

163 *Flow cytometry*

164 Samples were thawed rapidly at 37°C and washed twice in PBS containing 2% FCS and 2mM EDTA
165 by centrifuging at $400 \times g$ for 5 minutes. Cells were counted using a haemocytometer and approximately
166 300,000 PBMCs were added to tubes for incubation with fluorescently conjugated antibodies to identify
167 specific lymphocyte populations using eight-colour flow cytometry (FACS-CANTO, Becton–Dickinson, San
168 Jose, USA). The following monoclonal antibodies (mAbs) were used: anti-CD45RA-FITC clone # HI100,
169 anti-CD197 (CCR7)-PE clone # 150503, anti-CD56-PE-Cy7 clone # B159, anti-CD8-APC clone # RPA-T8,
170 anti-CD3-APC-Cy7 clone # SK7 (BD Pharmingen, San Diego, USA), anti-CLA-V450 clone # HECA-452
171 (BioLegend, San Diego, USA) and anti-CD16-V500 clone # 3G8 (BD Horizon, San Diego, USA). In addition,
172 7-aminoactinomycin D (7-AAD; PerCP channel; BD Pharmingen, San Diego, USA) was used to exclude
173 necrotic and apoptotic cells. For validation purposes, fluorescence-minus-one (FMO) tubes were used on
174 separate samples from three healthy donors to establish negative and positive gating strategies for CLA
175 expression.

176 Flow cytometry data were analysed using FlowJo version 7.6.5 (FlowJo LLC, Oregon, USA). Briefly,
177 lymphocytes were gated on the forward *versus* sideways scatter, and 7AAD $^{-}$ cells were divided into CD3 $^{+}$ or
178 CD3 $^{-}$ cells. Cytotoxic T cells were identified as being CD3 $^{+}$ CD8 $^{+}$ and further differentiated into naïve
179 (‘NA’; CCR7 $^{+}$ CD45RA $^{+}$), central memory (‘CM’; CCR7 $^{+}$ CD45RA $^{-}$), effector memory (‘EM’;
180 CCR7 $^{-}$ CD45RA $^{-}$) or CD45RA $^{+}$ effector memory cells (‘EMRA’; CCR7 $^{-}$ CD45RA $^{+}$). CD3 $^{-}$ cells were
181 further differentiated into CD56 $^{\text{dim}}$ ‘cytotoxic’ NK cells or CD56 $^{\text{high}}$ ‘regulatory’ NK cells, as proposed by
182 Cooper and colleagues (2001), using CD16 and CD56 dotplot gating (8). Within each subpopulation identified
183 above, cells were subdivided based on their expression of CLA (CLA $^{+}$ or CLA $^{-}$) in SSC (side scatter) versus
184 V450-CLA mode. Absolute cell counts for each subpopulation were computed by multiplying the percentage
185 of cells within the CD3 $^{+}$ or CD3 $^{-}$ gates by the total lymphocyte count (obtained via the Coulter principle).

186 This value and subsequent absolute subpopulation counts were multiplied by percentage values for gated
187 daughter subpopulations.

188

189 *Statistical analyses*

190 All results are presented as mean \pm standard deviation unless otherwise stated. Statistical calculations
191 were performed on IBM SPSS for Windows Version 21. Kolmogorov-Smirnov tests confirmed that all data
192 were normally distributed at all time points and for all variables. Repeated Measures Analysis of Variance
193 (ANOVA) tests were used to contrast changes over time (baseline, exercise and 30 min post-exercise), and
194 between the two exercise conditions (HIIE and continuous exercise) for each cell type. Within-trial main
195 effects of time for each cell type were calculated in separate individual ANOVAs. Post-hoc pairwise
196 comparisons were made with Bonferroni adjustments for multiple comparisons. Between trial differences at a
197 given time point were assessed by paired *t*-tests. Statistical significance was accepted at the $p < .05$ level.

RESULTS

All participants completed both the continuous exercise and HIIE tasks. Total energy expenditure was significantly higher after continuous exercise compared to HIIE (HIIE: 2.69 ± 0.31 Kcal/kg; continuous exercise: 3.54 ± 0.48 Kcal/kg; $F_{(1,17)} = 20.0$, $p < .05$). Similarly, RPE_{peak} was significantly higher after continuous exercise compared to HIIE (HIIE: 16 ± 2 (Borg scale); continuous exercise: 18 ± 1 (Borg scale); $F_{(1,17)} = 4.6$, $p < .05$). Workload was higher after HIIE compared to continuous exercise (HIIE: 2.74 ± 0.54 Watts/kg; continuous exercise: 2.18 ± 0.33 Watts/kg; $F_{(1,17)} = 7.1$, $p < .05$). No differences were found between trials for HR_{peak} (HIIE: 178 ± 14 beats/min; continuous exercise: 187 ± 7 beats/min; $p > .05$).

Effects of HIIE and continuous exercise on the number of total leukocytes and leukocyte subpopulations

Total leukocyte counts increased ($p < .05$) from baseline after both the HIIE (5.31 ± 1.15 [$\times 10^9/L$] to 8.92 ± 2.77) and continuous exercise (5.57 ± 0.94 to 10.73 ± 2.14) conditions. Similarly, total lymphocyte counts also increased ($p < .05$) from baseline after both the HIIE (1.83 ± 0.47 to 3.64 ± 1.46) and continuous exercise (1.91 ± 0.41 to 4.83 ± 1.33) conditions, with a larger mean change observed during continuous exercise, compared to HIIE, but these differences were not significant ($p > .05$). 30 min after each exercise condition, lymphocyte numbers between trials were similar (HIIE: 1.67 ± 0.37 ; continuous exercise: 1.84 ± 0.35) and had returned to pre-exercise levels ($p > .05$, compared to baseline).

Effects of HIIE and continuous exercise on the number of total CD3+ T cells and CD3+CD8+ T cells

Table 1 displays total CD3+ T cells and CD3+CD8+ T cell numbers during both exercise tasks. Total CD3+ T cells increased significantly during both conditions ($p < .01$), with larger mean increases during continuous exercise, compared to HIIE, but these between trial differences were not significant ($p > .05$). 30 min after each condition, CD3+ T cell numbers between trials were similar and had returned to pre-exercise levels. CD3+CD8+ T cells increased significantly during both conditions ($p < .05$), with larger increases during continuous exercise compared to HIIE; again, these differences between trials were not significant ($p > .05$). 30 min after each condition, CD3+CD8+ numbers between trials were similar and had returned to pre-

exercise levels. Replicating previous findings (6, 39), CD3+CD8+ T cell subsets were mobilised in a stepwise manner, with CD3+CD8+ EMRA T cells mobilising more than EM, CM and NA cells. Within these subsets, there were no significant differences in the magnitude of mobilisation between continuous exercise or HIIE trials.

228

Effects of HIIE and continuous exercise on the numbers of CLA+ and CLA- CD3+ T cells and CD3+CD8+ T cells

Approximately 14% of total CD3+ T cells were CLA+ at baseline, and despite an increase in the number of CLA+ CD3+ T cells during both exercise conditions ($p < .05$), the proportion of CLA+ CD3+ T cells decreased slightly during exercise due to a larger influx of CLA- CD3+ T cells ($p = .05$) (see Table 1). At 30 minutes post-exercise the number and proportion of CLA+ CD3+ T cells returned to pre-exercise values.

Approximately 9% of CD3+CD8+ T cells were CLA+ at baseline, and, in a similar manner to total CD3+ T cells, the number of CLA+ CD3+CD8+ T cells increased during both exercise conditions ($p < .05$), before returning to baseline levels at 30 minutes post-exercise (see Table 1). However, there was no change ($p > .05$) in the proportion of CLA+ CD3+CD8+ T cells during exercise. No differences were observed in the magnitude of mobilisation between CLA+ and CLA- cells in any of these broad T cell subsets between HIIE and continuous exercise.

242

Effects of HIIE and continuous exercise on the numbers of CLA+ and CLA- CD3+CD8+ T cell subsets

In accordance with prior literature (7, 31), very few of the CD3+CD8+ NA cell population expressed CLA (median = 1.35%) and the few CLA+ naïve cells did not change during exercise (data not shown). The largest proportion of CLA+ CD3+CD8+ T cells were of the CM (~20%) and EM (~23%) phenotypes, with only a small fraction of EMRA cells expressing CLA (<7%). As illustrated in Figure 1, the percentage change in the number of CLA+ cells increases in a stepwise mobilisation pattern (i.e., EMRA > EM > CM) in

response to both exercise conditions, and this pattern is also evident in CLA⁻ cells. However, as illustrated in Figure 2, the proportion of CLA⁺ cells per memory CD3⁺CD8⁺ T cell subset actually shows a marginal and significant ($p < 0.05$) decline during exercise, before returning to baseline levels at 30 min post-exercise. Thus, CLA⁺ cells do not appear to be major contributors to the large influx of effector cells observed in the peripheral bloodstream during exercise; this is illustrated by a greater mobilisation of CLA⁻, compared to CLA⁺, EMRA CD3⁺CD8⁺ T cells in Figure 1.

Effects of HIIE and continuous exercise on the numbers of NK cells, and CLA⁺ and CLA⁻ cells within the NK cell subsets

Table 2 displays changes in the numbers of CD3⁺CD56⁺ NK cells in response to continuous exercise and HIIE. As expected, NK cells were extremely sensitive to exercise stress, exhibiting significant differences between the two exercise modalities ($p < .05$), with HIIE resulting in a ~400% increase and continuous exercise resulting in a ~600% change during exercise. With regards to the major CD56⁺ NK cell subsets, CD56^{dim} ‘cytotoxic’ cells were the most sensitive to exercise stress, with HIIE resulting in a 550% increase and continuous exercise resulting in a 725% increase ($p < .05$). As expected, CD56^{bright} ‘regulatory’ NK cells were less sensitive to exercise stress and no differences were found between exercise conditions ($p > .05$); HIIE resulted in a 100% increase and continuous exercise resulted in a 200% increase.

At baseline, approximately ~22% of CD3⁺CD56⁺ cells were CLA⁺, with a higher proportion of CD56^{bright} ‘regulatory’ NK cells positive for CLA (~67%) than CD56^{dim} ‘cytotoxic’ NK cells (~17.5%). During exercise, the total number of CLA⁺ NK cells increased (HIIE: ~230%; continuous exercise: ~350%), and, as expected, this was driven by a larger increase in CLA⁺ CD56^{dim} cells compared to the CLA⁺ CD56^{bright} cells. As a consequence of a greater influx of CLA⁻ NK cells (HIIE: ~500%; continuous exercise: ~650% compared to baseline), primarily comprised of CLA⁻CD56^{dim} cells during exercise (Figure 3), the proportion of CLA⁺ cells in the total NK cell pool was actually reduced during exercise (Figure 4). Thus, CLA⁺ NK cells do not appear to be major contributors to the large influx of effector cells observed in the peripheral bloodstream during exercise.

275 **DISCUSSION**

276 This study assessed the mobilisation propensity of CLA⁺ CD8⁺ T cells into the peripheral
277 bloodstream in response to intensive exercise. A preferential mobilisation of this cell phenotype would fulfil a
278 component of the stress redistribution theory (10), which hypothesises that effector memory cells
279 preferentially mobilised by exercise have a phenotype that enables post-exercise migration to cutaneous sites
280 of wound healing or active inflammation (e.g., injured skin). We examined the number of CLA⁺ and CLA⁻
281 cells because CLA avidly binds CD62E; CD62E is thought to be the primary initiator of routine CD8⁺ T cell
282 transmigration to the skin, and is also central to T cell tethering to cytokine-activated endothelium at inflamed
283 cutaneous sites (14). We found that exercise resulted in an increase in the number of memory CLA⁺
284 CD3⁺CD8⁺ T cells. However, the numerical contribution of CLA⁺ memory cells to exercise-induced
285 lymphocytosis was inferior to CLA⁻ cells, and, as a consequence, the proportion of CLA⁺ cells amongst the
286 total memory CD8⁺ T cell pool showed a decline during exercise. Furthermore, the most exercise-sensitive
287 subset (EMRA CD8⁺ T cells) were largely CLA⁻, suggesting that the most exercise-responsive T cell subset
288 mobilised by exercise do not have a phenotype that would enable rapid transmigration to sites of active
289 cutaneous inflammation.

290 Until now, the effect of exercise on the mobilisation of CLA⁺ cells has not yet been investigated. A
291 study investigating the effects of acute psychological stress on circulating CLA⁺ CD8⁺ cells found a marginal
292 decline in the number of CLA⁺ CD8⁺ cells during the stressor (2). These aforementioned results differed to
293 the findings of our study, as we observed an approximate doubling in the number of CLA⁺ CD3⁺CD8⁺ cells
294 during exercise. It was proposed by Atanackovic et al. (2006) that CLA⁺ CD8⁺ T cells had already initiated
295 transmigration to sites of inflammation during acute stress. However, evidence exists to suggest that elevated
296 levels of epinephrine selectively decreases adhesion of CD8⁺ T cells (12) and NK cells (3) to endothelial
297 cells, and may thus be a contributing mechanism for the maintenance of CLA⁺ cells in the peripheral
298 bloodstream, as observed in our exercise trials. We extended our CLA⁺ analyses to four distinct CD8⁺ T cell
299 subsets conventionally identified using the cell-surface markers CCR7 and CD45RA (17). In agreement with
300 prior literature showing that the majority of CLA⁺ CD3⁺ cells are CD45RO⁺ (7), we found that few naïve
301 (CCR7⁺CD45RA⁺) CD8⁺ T cells expressed CLA (31). This did not change in response to exercise. In further

302 agreement with Clark et al. (2006), we found a majority of CLA⁺ cells were of the central memory (CCR7⁺
 303 CD45RA⁻) phenotype and fewer were effector memory (CCR7⁻CD45RA⁻) CD8⁺ T cells; these CLA⁺
 304 effector memory cells showed the largest increase during exercise, though the numerical increase was small.

305 On one hand, the findings of this study are supportive of the stress redistribution model in that
 306 exercise evoked an increase in the total number of peripheral blood CLA⁺ CD8⁺ T cells, which are
 307 phenotypically consistent with skin-homing T cells. On the other hand, the absolute number of CLA⁺ cells
 308 mobilised was small, and the relative magnitude of mobilisation was much less than CLA⁻ CD8⁺ cells.

309 Furthermore, few EMRA CD8⁺ T cells – conventionally the most exercise-sensitive CD8⁺ subset – expressed
 310 CLA, and did not increase in number as much as CLA⁻ EMRA CD8⁺ T cells. Thus, the CD8⁺ T cells with
 311 the greatest propensity for exercise-induced mobilisation and extravasation are CLA⁻ and do not exhibit a
 312 phenotype characteristic for rapid homing to cutaneous sites or sites of endothelial inflammation. Given that
 313 the large majority of CLA⁺ cells reside in the skin (7), it is possible to conclude that the skin is not a major
 314 contributor/source of cells to exercise-induced lymphocytosis. The findings of this study are not surprising:
 315 CLA⁺ cells are not as susceptible to immunosenescence (25), and are thus strikingly different to the most
 316 exercise-sensitive cells, which have shorter telomeres, high expression of CD57 (35, 36); and exhibit
 317 exaggerated exercise-induced mobilisation in participants seropositive for cytomegalovirus (39). In addition, it
 318 has been demonstrated in rodents that exercise redeploys large numbers of T cells to the lungs (20). Lung-
 319 derived T cells are all CLA⁻, but positive for other homing molecules including CCR5 and CXCR3 (5).

320 Similarly, exercise also redeploys T cells to the Peyer's patches and bone marrow (19, 20) and this is likely
 321 governed by other site-specific homing molecules like LPAM-1 (37) or VLA-4 (23, 26) rather than CLA. This
 322 allocation of certain cells to defined parts of the body might represent a homeostatic immune-surveillance
 323 response (10), or, instead, it has been hypothesised that senescent T cells are mobilised into the blood to
 324 facilitate their subsequent apoptosis in peripheral tissues (21, 34), which may contribute to progenitor cell
 325 mobilisation after exercise (24). With regards to cutaneous surveillance against tumours and stressed tissue
 326 cells, it is unlikely this is tasked by $\alpha\beta$ CD8⁺ T cells alone (13), and is likely supported by $\gamma\delta$ T cells and NK
 327 cells, which are highly responsive to acute stress (1, 6), and have the migratory capacity to enter cutaneous
 328 sites (13). We found that CLA⁺ CD56⁺ NK cells were substantially mobilised by exercise, but, much like
 329 CD8⁺ T cells, were outnumbered by the mobilisation of CLA⁻ NK cells.

330 This is the first study to investigate and compare the effects of HIIE (also referred to as HIIT when
331 repeated over several weeks or months) and continuous exercise on T cell and NK cell subset mobilisation
332 responses. HIIE typically involves shorter and more intense bouts of exercise than more traditional forms
333 (e.g., 30 minutes of moderate intensity running or cycling) and is thought to be more attractive and better
334 tolerated by participants. HIIE has received considerable recent attention as an effective means of achieving
335 certain physiological adaptations (e.g., improved insulin sensitivity and cardiorespiratory fitness, but probably
336 not weight loss) in healthy and diseased populations that are similar or superior to traditional endurance-based
337 exercise (16).

338 Although we found a trend whereby a greater mobilisation of lymphocytes was observed during
339 continuous exercise compared to HIIE, these differences were not significant for the majority of lymphocyte
340 subpopulations investigated. We found that CD56 NK cells – conventionally the most exercise responsive
341 cells – were mobilised to a greater extent during continuous exercise compared to HIIE, a response that was
342 driven by larger increases to CD56^{dim} NK cells. It may be that similar intensity-dependent effects would have
343 been found for CD8⁺ T cells, and other subsets, if a larger group of participants had been tested.

344 A limitation of this study was that post-exercise blood samples were collected 30 minutes after
345 exercise cessation, rather than after 60 minutes, when NK and CD8⁺ T cell lymphocytopenia is more
346 pronounced (6, 39). This was a consequence of practical and logistical constraints imposed by the broader
347 study that was being undertaken (41). At 30 min post-exercise, we found that all cell phenotypes were present
348 in peripheral blood in similar numbers to pre-exercise levels. It would be of interest to investigate the
349 contribution of CLA⁺ T cells to lymphocytopenia at later time points. An additional practical limitation was
350 the cryopreservation of PBMCs, which may disproportionately affect the viability of some cell phenotypes –
351 however this is unlikely to affect the within-subject model used in our study. A further consideration for
352 future studies surrounds the ongoing debate over the optimal phenotyping of skin-homing CD8⁺ T cells. In
353 addition to CLA, CCR4 and CCR8 may be useful in differentiating between CD8⁺ T cells involved in normal
354 cutaneous immune-surveillance to those involved in acute or chronic inflammation (7, 33). Finally, our
355 investigations were limited to healthy individuals with no apparent cutaneous inflammation. Investigation of
356 exercise-induced lymphocyte skin-homing in experimental models of infection or in clinical populations

including psoriasis is warranted, and more invasive techniques such as biopsy may provide further insights into exercise-induced homing to cutaneous or inflammatory sites.

CONCLUSION

A greater number of CLA⁻ T cells and NK cells were mobilised into peripheral blood than CLA⁺ counterpart T cells and NK cells during exercise. Furthermore, the majority of EMRA T cells and CD56^{dim} cells – i.e., conventionally the most exercise-responsive cells – did not express CLA. Together, these findings demonstrate that CLA⁺ cells are not major contributors to exercise lymphocytosis, thus providing preliminary evidence that the skin is not a major origin, or homing-destination, of exercise-sensitive lymphocytes. We conclude that the most exercise-sensitive lymphocytes likely migrate from, and to, non-cutaneous sites post-exercise.

ACKNOWLEDGEMENTS: The results of this present study do not constitute endorsement by ACSM. The flow cytometric analyses for this study were funded by the University of Birmingham Clinical Immunology Service.

CONFLICTS OF INTEREST: None declared

374 REFERENCES

- 375 1. Anane LH, Edwards KM, Burns VE et al. Mobilization of gammadelta T lymphocytes in response to
376 psychological stress, exercise, and beta-agonist infusion. *Brain, behavior, and immunity*.
377 2009;23(6):823-9.
- 378 2. Atanackovic D, Schnee B, Schuch G et al. Acute psychological stress alerts the adaptive immune
379 response: stress-induced mobilization of effector T cells. *Journal of neuroimmunology*. 2006;176(1-
380 2):141-52.
- 381 3. Benschop RJ, Nijkamp FP, Ballieux RE, Heijnen CJ. The effects of beta-adrenoceptor stimulation on
382 adhesion of human natural killer cells to cultured endothelium. *British journal of pharmacology*.
383 1994;113(4):1311-6.
- 384 4. Berg EL, Yoshino T, Rott LS et al. The cutaneous lymphocyte antigen is a skin lymphocyte homing
385 receptor for the vascular lectin endothelial cell-leukocyte adhesion molecule 1. *The Journal of*
386 *experimental medicine*. 1991;174(6):1461-6.
- 387 5. Campbell JJ, Brightling CE, Symon FA et al. Expression of chemokine receptors by lung T cells from
388 normal and asthmatic subjects. *J Immunol*. 2001;166(4):2842-8.
- 389 6. Campbell JP, Riddell NE, Burns VE et al. Acute exercise mobilises CD8+ T lymphocytes exhibiting an
390 effector-memory phenotype. *Brain, behavior, and immunity*. 2009;23(6):767-75.
- 391 7. Clark RA, Chong B, Mirchandani N et al. The vast majority of CLA+ T cells are resident in normal skin.
392 *J Immunol*. 2006;176(7):4431-9.
- 393 8. Cooper MA, Fehniger TA, Caligiuri MA. The biology of human natural killer-cell subsets. *Trends in*
394 *immunology*. 2001;22(11):633-40.
- 395 9. Dhabhar FS. Psychological stress and immunoprotection versus immunopathology in the skin. *Clinics*
396 *in dermatology*. 2013;31(1):18-30.
- 397 10. Dhabhar FS, Malarkey WB, Neri E, McEwen BS. Stress-induced redistribution of immune cells--from
398 barracks to boulevards to battlefields: a tale of three hormones--Curt Richter Award winner.
399 *Psychoneuroendocrinology*. 2012;37(9):1345-68.
- 400 11. Dimitroff CJ, Kupper TS, Sackstein R. Prevention of leukocyte migration to inflamed skin with a novel
401 fluorosugar modifier of cutaneous lymphocyte-associated antigen. *The Journal of clinical*
402 *investigation*. 2003;112(7):1008-18.
- 403 12. Dimitrov S, Lange T, Born J. Selective mobilization of cytotoxic leukocytes by epinephrine. *J Immunol*.
404 2010;184(1):503-11.
- 405 13. Ebert LM, Meuter S, Moser B. Homing and function of human skin gammadelta T cells and NK cells:
406 relevance for tumor surveillance. *J Immunol*. 2006;176(7):4331-6.
- 407 14. Fuhlbrigge RC, Kieffer JD, Armerding D, Kupper TS. Cutaneous lymphocyte antigen is a specialized
408 form of PSGL-1 expressed on skin-homing T cells. *Nature*. 1997;389(6654):978-81.
- 409 15. Gannon GA, Rhind SG, Shek PN, Shephard RJ. Differential cell adhesion molecule expression and
410 lymphocyte mobilisation during prolonged aerobic exercise. *European journal of applied physiology*.
411 2001;84(4):272-82.
- 412 16. Gibala MJ, Little JP, Macdonald MJ, Hawley JA. Physiological adaptations to low-volume, high-
413 intensity interval training in health and disease. *The Journal of physiology*. 2012;590(Pt 5):1077-84.
- 414 17. Hamann D, Baars PA, Rep MH et al. Phenotypic and functional separation of memory and effector
415 human CD8+ T cells. *The Journal of experimental medicine*. 1997;186(9):1407-18.
- 416 18. Hoshino T, Yamada A, Honda J et al. Tissue-specific distribution and age-dependent increase of
417 human CD11b+ T cells. *J Immunol*. 1993;151(4):2237-46.
- 418 19. Kruger K, Lechtermann A, Fobker M, Volker K, Mooren FC. Exercise-induced redistribution of T
419 lymphocytes is regulated by adrenergic mechanisms. *Brain, behavior, and immunity*. 2008;22(3):324-
420 38.
- 421 20. Kruger K, Mooren FC. T cell homing and exercise. *Exercise immunology review*. 2007;13:37-54.
- 422 21. Kruger K, Mooren FC. Exercise-induced leukocyte apoptosis. *Exercise immunology review*.
423 2014;20:117-34.

- 424 22. Kurokawa Y, Shinkai S, Torii J, Hino S, Shek PN. Exercise-induced changes in the expression of surface
425 adhesion molecules on circulating granulocytes and lymphocytes subpopulations. *European journal*
426 *of applied physiology and occupational physiology*. 1995;71(2-3):245-52.
- 427 23. Mazo IB, Honczarenko M, Leung H et al. Bone marrow is a major reservoir and site of recruitment for
428 central memory CD8+ T cells. *Immunity*. 2005;22(2):259-70.
- 429 24. Mooren FC, Kruger K. Apoptotic lymphocytes induce progenitor cell mobilization after exercise. *J*
430 *Appl Physiol (1985)*. 2015:jap 00287 2015.
- 431 25. Neuber K, Schmidt S, Mensch A. Telomere length measurement and determination of
432 immunosenescence-related markers (CD28, CD45RO, CD45RA, interferon-gamma and interleukin-4)
433 in skin-homing T cells expressing the cutaneous lymphocyte antigen: indication of a non-ageing T-cell
434 subset. *Immunology*. 2003;109(1):24-31.
- 435 26. Papayannopoulou T, Priestley GV, Nakamoto B, Zafiropoulos V, Scott LM. Molecular pathways in
436 bone marrow homing: dominant role of alpha(4)beta(1) over beta(2)-integrins and selectins. *Blood*.
437 2001;98(8):2403-11.
- 438 27. Picker LJ, Kishimoto TK, Smith CW, Warnock RA, Butcher EC. ELAM-1 is an adhesion molecule for
439 skin-homing T cells. *Nature*. 1991;349(6312):796-9.
- 440 28. Picker LJ, Michie SA, Rott LS, Butcher EC. A unique phenotype of skin-associated lymphocytes in
441 humans. Preferential expression of the HECA-452 epitope by benign and malignant T cells at
442 cutaneous sites. *The American journal of pathology*. 1990;136(5):1053-68.
- 443 29. Picker LJ, Treer JR, Ferguson-Darnell B, Collins PA, Bergstresser PR, Terstappen LW. Control of
444 lymphocyte recirculation in man. II. Differential regulation of the cutaneous lymphocyte-associated
445 antigen, a tissue-selective homing receptor for skin-homing T cells. *J Immunol*. 1993;150(3):1122-36.
- 446 30. Robert C, Kupper TS. Inflammatory skin diseases, T cells, and immune surveillance. *The New England*
447 *journal of medicine*. 1999;341(24):1817-28.
- 448 31. Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with
449 distinct homing potentials and effector functions. *Nature*. 1999;401(6754):708-12.
- 450 32. Santamaria Babi LF, Moser R, Perez Soler MT, Picker LJ, Blaser K, Hauser C. Migration of skin-homing
451 T cells across cytokine-activated human endothelial cell layers involves interaction of the cutaneous
452 lymphocyte-associated antigen (CLA), the very late antigen-4 (VLA-4), and the lymphocyte function-
453 associated antigen-1 (LFA-1). *J Immunol*. 1995;154(4):1543-50.
- 454 33. Schaerli P, Ebert LM, Moser B. Comment on "The vast majority of CLA+ T cells are resident in normal
455 skin". *J Immunol*. 2006;177(3):1375-6; author reply 6-7.
- 456 34. Simpson RJ. Aging, persistent viral infections, and immunosenescence: can exercise "make space"?
457 *Exercise and sport sciences reviews*. 2011;39(1):23-33.
- 458 35. Simpson RJ, Cosgrove C, Chee MM et al. Senescent phenotypes and telomere lengths of peripheral
459 blood T-cells mobilized by acute exercise in humans. *Exercise immunology review*. 2010;16:40-55.
- 460 36. Simpson RJ, Cosgrove C, Ingram LA et al. Senescent T-lymphocytes are mobilised into the peripheral
461 blood compartment in young and older humans after exhaustive exercise. *Brain, behavior, and*
462 *immunity*. 2008;22(4):544-51.
- 463 37. Strauch UG, Lifka A, Gossler U, Kilshaw PJ, Clements J, Holzmann B. Distinct binding specificities of
464 integrins alpha 4 beta 7 (LPAM-1), alpha 4 beta 1 (VLA-4), and alpha IEL beta 7. *International*
465 *immunology*. 1994;6(2):263-75.
- 466 38. Thatte J, Dabak V, Williams MB, Braciale TJ, Ley K. LFA-1 is required for retention of effector CD8 T
467 cells in mouse lungs. *Blood*. 2003;101(12):4916-22.
- 468 39. Turner JE, Aldred S, Witard OC, Drayson MT, Moss PM, Bosch JA. Latent cytomegalovirus infection
469 amplifies CD8 T-lymphocyte mobilisation and egress in response to exercise. *Brain, behavior, and*
470 *immunity*. 2010;24(8):1362-70.
- 471 40. Viswanathan K, Dhabhar FS. Stress-induced enhancement of leukocyte trafficking into sites of
472 surgery or immune activation. *Proceedings of the National Academy of Sciences of the United States*
473 *of America*. 2005;102(16):5808-13.

- 474 41. Wadley AJ, Chen YW, Bennett SJ et al. Monitoring changes in Thioredoxin and over-oxidised
475 Peroxiredoxin in response to exercise in humans. *Free radical research*. 2015;49(3):290-8.
- 476 42. Wadley AJ, Chen YW, Lip GY, Fisher JP, Aldred S. Low volume-high intensity interval exercise elicits
477 antioxidant and anti-inflammatory effects in humans. *Journal of sports sciences*. 2015;27:1-9.
- 478 43. Williams MB, Butcher EC. Homing of naive and memory T lymphocyte subsets to Peyer's patches,
479 lymph nodes, and spleen. *J Immunol*. 1997;159(4):1746-52.

480

481

482

483

484

FIGURE CAPTIONS

Figure 1. Mean \pm SEM changes in the proportion of CD8⁺ T cell subsets (CM = central memory; EM = effector memory; EMRA = CD45RA⁺ effector memory) that were either positive or negative for cell surface CLA, at baseline and immediately after Continuous exercise. It was found that CLA⁺ CD45RA⁺ effector memory CD8⁺ T cells were mobilised to a greater magnitude than CLA⁻ CD45RA⁺ effector memory CD8⁺ T cells; similar patterns were found for HIIE (data not shown).

* Indicates significant difference between CLA⁺ and CLA⁻ ($p < .05$).

Figure 2. Mean \pm SEM changes in the proportion of CD8⁺ T cell subsets (NA = naïve; CM = central memory; EM = effector memory; EMRA = CD45RA⁺ effector memory) that were positive for CLA, from pre- to immediately post-continuous exercise. A reduction in the proportion of CLA⁺ CM, EM and EMRA cells was observed immediately after continuous exercise compared to baseline; similar patterns were observed during HIIE (data not shown).

* Indicates significant differences in the proportion of CLA⁺ cells between pre- and post-continuous exercise ($p < .05$).

Figure 3. Mean \pm SEM changes in the proportion of NK cell subsets that were either CLA⁺ or CLA⁻, at baseline and immediately after continuous exercise. A trend was observed whereby CLA⁻ CD56^{dim} cells were mobilised to a greater extent than CLA⁺ CD56^{dim} NK cells ($p = .058$); similar patterns were found for HIIE (data not shown).

Figure 4. Mean \pm SEM changes in the proportion of NK cell subsets that were CLA⁺, from pre- to immediately post-continuous exercise. A reduction in the proportion of CLA⁺ CD56^{bright} and CD56^{dim} NK cells was observed from pre- to post-exercise; similar patterns were observed during HIIE (data not shown).

* Indicates significant differences in the proportion of CLA⁺ cells between pre- and post-continuous exercise ($p < .05$).

Table 1. Changes in the numbers of T-lymphocytes and T-lymphocyte subpopulations in response to HIIE and continuous exercise (mean \pm SD).

Cells per μ L	Condition	Baseline	Exercise	+30min	Main effects of time	Time \times task interaction
CD3+	HIIE	1494.4 \pm 406.2	2508.8 \pm 1076.5*	1330.2 \pm 293.2	$F_{(2,16)} = 16.18; p = .002$	$F_{(2,32)} = 2.06; p = \text{NS}$
	Continuous Ex	1454.1 \pm 321.9	3049.6 \pm 794.4***	1416.3 \pm 363.1	$F_{(2,16)} = 42.97; p < .001$	
CD3+ CLA+	HIIE	189.2 \pm 89.7	316.2 \pm 237.0	186.8 \pm 95.7	$F_{(2,16)} = 6.80; p = .030$	$F_{(2,32)} = .37; p = \text{NS}$
	Continuous Ex	183.0 \pm 91.4	350.6 \pm 166.6**	196.7 \pm 123.2	$F_{(2,16)} = 26.06; p < .001$	
CD3+ CLA-	HIIE	1305.2 \pm 376.2	2192.6 \pm 894.2**	1143.4 \pm 260.2	$F_{(2,16)} = 17.72; p = .002$	$F_{(2,32)} = 2.33; p = \text{NS}$
	Continuous Ex	1271.0 \pm 293.3	2699.0 \pm 276.0***	1219.6 \pm 291.7	$F_{(2,16)} = 41.40; p < .001$	
CD3+ CD8+	HIIE	509.5 \pm 181.9	1000.9 \pm 654.7	463.1 \pm 174.7	$F_{(2,16)} = 9.46; p = .014$	$F_{(2,32)} = .89; p = \text{NS}$
	Continuous Ex	491.2 \pm 162.3	1185.0 \pm 398.8***	490.4 \pm 195.4	$F_{(2,16)} = 48.99; p < .001$	
CD3+ CD8+ CLA+	HIIE	41.3 \pm 21.6	77.9 \pm 65.6	43.2 \pm 29.3	$F_{(2,16)} = 6.22; p = .035$	$F_{(2,32)} = .28; p = \text{NS}$
	Continuous Ex	40.7 \pm 26.2	86.9 \pm 53.8**	43.8 \pm 32.7	$F_{(2,16)} = 23.95; p = .001$	
CD3+ CD8+ CLA-	HIIE	468.2 \pm 163.4	923.0 \pm 591.5	420.0 \pm 148.7	$F_{(2,16)} = 9.76; p = .013$	$F_{(2,32)} = .95; p = \text{NS}$
	Continuous Ex	450.4 \pm 141.2	1098.1 \pm 355.0***	446.5 \pm 166.9	$F_{(2,16)} = 48.96; p < .001$	
Naïve CD8+	HIIE	239.9 \pm 93.1	341.6 \pm 144.1*	192.8 \pm 62.2	$F_{(2,16)} = 14.14; p < .001$	$F_{(2,32)} = 1.51; p = \text{NS}$
	Continuous Ex	215.9 \pm 76.1	377.9 \pm 69.0***	198.5 \pm 56.4	$F_{(2,16)} = 55.18; p < .001$	
Naïve CD8+ CLA+	HIIE	3.4 \pm 1.4	5.2 \pm 2.6	3.0 \pm 1.4	$F_{(2,16)} = 10.20; p = .001$	$F_{(2,32)} = .97; p = \text{NS}$
	Continuous Ex	3.5 \pm 1.8	6.3 \pm 3.4*	3.5 \pm 2.2	$F_{(2,16)} = 17.67; p = .002$	
Naïve CD8+ CLA-	HIIE	236.5 \pm 92.2	336.4 \pm 142.3*	189.8 \pm 62.0	$F_{(2,16)} = 14.13; p < .001$	$F_{(2,32)} = 1.49; p = \text{NS}$
	Continuous Ex	212.4 \pm 75.8	370.5 \pm 70.3***	195.0 \pm 55.4	$F_{(2,16)} = 54.09; p < .001$	
CM	HIIE	118.8 \pm 37.2	221.8 \pm 97.1*	124.7 \pm 55.9	$F_{(2,16)} = 21.50; p = .001$	$F_{(2,32)} = 3.85; p = \text{NS}$
	Continuous Ex	121.9 \pm 64.2	278.6 \pm 96.2***	132.3 \pm 85.4	$F_{(2,16)} = 101.54; p < .001$	
CM CD8+ CLA+	HIIE	20.2 \pm 8.3	35.6 \pm 25.0	21.7 \pm 12.7	$F_{(2,16)} = 7.01; p = .026$	$F_{(2,32)} = .29; p = \text{NS}$
	Continuous Ex	20.0 \pm 11.8	38.9 \pm 20.7**	21.3 \pm 14.7	$F_{(2,16)} = 25.16; p < .001$	
CM CD8+ CLA-	HIIE	98.6 \pm 30.4	186.2 \pm 72.9**	103.1 \pm 44.2	$F_{(2,16)} = 27.81; p < .001$	$F_{(2,32)} = 5.69; p = .017$
	Continuous Ex	101.9 \pm 53.5	239.7 \pm 77.4***	110.6 \pm 71.8	$F_{(2,16)} = 123.76; p < .001$	
EM	HIIE	64.4 \pm 29.9	144.9 \pm 70.8**	69.5 \pm 37.2	$F_{(2,16)} = 26.74; p = .001$	$F_{(2,32)} = 3.30; p = \text{NS}$
	Continuous Ex	66.6 \pm 39.3	184.8 \pm 71.0***	75.6 \pm 52.0	$F_{(2,16)} = 80.73; p < .001$	
EM CD8+ CLA+	HIIE	12.3 \pm 10.5	24.7 \pm 27.5	13.7 \pm 13.4	$F_{(2,16)} = 4.73; p = \text{NS}$	$F_{(2,32)} = .13; p = \text{NS}$
	Continuous Ex	12.0 \pm 10.8	26.9 \pm 24.5*	13.6 \pm 14.0	$F_{(2,16)} = 11.22; p = .010$	
EM CD8+ CLA-	HIIE	52.1 \pm 20.5	120.2 \pm 44.7***	55.8 \pm 24.6	$F_{(2,16)} = 40.39; p < .001$	$F_{(2,32)} = 5.08; p = .026$
	Continuous Ex	54.6 \pm 29.7	157.8 \pm 51.3***	62.2 \pm 40.4	$F_{(2,16)} = 90.05; p < .001$	
EMRA	HIIE	88.3 \pm 87.0	292.6 \pm 390.3	76.1 \pm 69.7	$F_{(2,16)} = 4.07; p = \text{NS}$	$F_{(2,32)} = .13; p = \text{NS}$
	Continuous Ex	86.6 \pm 57.7	344.6 \pm 296.6*	83.8 \pm 52.1	$F_{(2,16)} = 9.16; p = .016$	
EMRA CD8+ CLA+	HIIE	5.4 \pm 4.0	12.4 \pm 12.6	4.9 \pm 3.7	$F_{(2,16)} = 5.35; p = .047$	$F_{(2,32)} = .29; p = \text{NS}$
	Continuous Ex	5.4 \pm 4.4	14.6 \pm 10.9**	5.4 \pm 4.6	$F_{(2,16)} = 17.13; p = .003$	
EMRA CD8+ CLA-	HIIE	80.9 \pm 84.0	280.2 \pm 378.7	71.2 \pm 66.8	$F_{(2,16)} = 4.02; p = \text{NS}$	$F_{(2,32)} = .12; p = \text{NS}$
	Continuous Ex	81.4 \pm 54.4	329.9 \pm 291.0*	78.4 \pm 48.6	$F_{(2,16)} = 8.75; p = .018$	

* $p < .05$ in comparison to baseline

** $p < .01$ in comparison to baseline

*** $p < .001$ in comparison to baseline

NS $p > .05$

Table 2. Changes in NK cell numbers in response to HIIE and continuous exercise (mean \pm SD).

Cells per μ L	Condition	Baseline	Exercise	+30min	Main effects of time	Time \times task interaction
CD56 ⁺	HIIE	112.4 \pm 44.8	591.9 \pm 361.0**	115.2 \pm 59.5	$F_{(2,16)} = 18.27; p = .003$	$F_{(2,32)} = 4.47; p = .05$
	Continuous Ex	164.3 \pm 144.2	1130.6 \pm 707.5**	157.6 \pm 159.9	$F_{(2,16)} = 22.64; p = .001$	
CD56 ⁺ CLA ⁺	HIIE	28.2 \pm 10.4	92.7 \pm 47.8**	29.2 \pm 12.4	$F_{(2,16)} = 23.96; p = .001$	$F_{(2,32)} = 3.37; p = \text{NS}$
	Continuous Ex	31.7 \pm 9.5	142.1 \pm 69.2**	31.9 \pm 10.9	$F_{(2,16)} = 26.10; p = .001$	
CD56 ⁺ CLA ⁻	HIIE	84.1 \pm 44.6	499.3 \pm 352.6*	86.0 \pm 56.5	$F_{(2,16)} = 14.13; p = .005$	$F_{(2,32)} = 3.75; p = \text{NS}$
	Continuous Ex	132.6 \pm 136.6	988.4 \pm 698.6**	125.7 \pm 153.2	$F_{(2,16)} = 18.14; p = .003$	
CD56 ^{bright}	HIIE	11.0 \pm 7.4	23.9 \pm 19.4*	15.2 \pm 13.0	$F_{(2,16)} = 9.98; p = .011$	$F_{(2,32)} = 2.17; p = \text{NS}$
	Continuous Ex	12.8 \pm 7.0	36.0 \pm 25.7*	15.6 \pm 7.8	$F_{(2,16)} = 11.05; p = .009$	
CD56 ^{bright} CLA ⁺	HIIE	7.6 \pm 5.5	15.0 \pm 12.7*	10.2 \pm 9.1	$F_{(2,16)} = 9.05; p = .015$	$F_{(2,32)} = 1.32; p = \text{NS}$
	Continuous Ex	8.4 \pm 4.7	20.3 \pm 15.8*	10.2 \pm 5.7	$F_{(2,16)} = 8.88; p = .015$	
CD56 ^{bright} CLA ⁻	HIIE	3.4 \pm 2.0	8.9 \pm 7.2*	5.0 \pm 4.0	$F_{(2,16)} = 10.14; p = .011$	$F_{(2,32)} = 1.79; p = \text{NS}$
	Continuous Ex	4.4 \pm 2.7	15.8 \pm 14.4	5.4 \pm 2.5	$F_{(2,16)} = 6.35; p = .035$	
CD56 ^{dim}	HIIE	101.4 \pm 44.5	568.0 \pm 343.6**	99.9 \pm 55.6	$F_{(2,16)} = 18.41; p = .003$	$F_{(2,32)} = 4.49; p = .05$
	Continuous Ex	151.5 \pm 47.2	1094.5 \pm 229.9**	141.9 \pm 53.1	$F_{(2,16)} = 22.72; p = .001$	
CD56 ^{dim} CLA ⁺	HIIE	20.6 \pm 10.0	77.7 \pm 43.8**	19.0 \pm 7.6	$F_{(2,16)} = 22.68; p = .001$	$F_{(2,32)} = 3.21; p = \text{NS}$
	Continuous Ex	23.3 \pm 7.8	121.8 \pm 60.8**	21.7 \pm 8.9	$F_{(2,16)} = 25.51; p = .001$	
CD56 ^{dim} CLA ⁻	HIIE	80.7 \pm 44.0	490.3 \pm 345.8*	81.0 \pm 55.0	$F_{(2,16)} = 14.16; p = .005$	$F_{(2,32)} = 3.75; p = \text{NS}$
	Continuous Ex	128.2 \pm 135.3	972.7 \pm 689.1**	120.3 \pm 152.3	$F_{(2,16)} = 18.19; p = .003$	

* $p < .05$ in comparison to baseline** $p < .01$ in comparison to baselineNS $p > .05$

Figure 1




Figure 2




Figure 3




Figure 4

